Rayaldee®
calcifediol ER capsules
30 mcg

Take Control of Your Prescription Costs

Learn about the Medicare Prescription Payment Plan (M3P)

DO YOU NEED HELP MANAGING YOUR PRESCRIPTION COSTS?

- Do you have **Medicare Part D coverage?**
- Do you have **high or unpredictable** prescription costs?
- Have you ever **delayed or skipped** a prescription because of cost?
- Would spreading your prescription costs over time make them **easier to manage?**

If you answered **YES** to any of these questions, the Medicare Prescription Payment Plan may be right for you.

WHAT IS THE MEDICARE PRESCRIPTION PAYMENT PLAN (M3P)?

The Medicare Prescription Payment Plan (M3P) is a program that allows people enrolled in Medicare Part D to spread out their out-of-pocket costs over the course of the calendar year for prescription drugs covered by their plan. This payment option is available through all Medicare health plans, participation is voluntary, and there is no cost to participate.[1]

If you choose M3P:

You pay **$0** at the pharmacy for prescriptions covered by your Medicare drug plan.[1]

Your health of drug plan will send you a **monthly bill** for your share of prescription costs.[1]

$2,400

From January through December, **you will not spend more than $2,400 total** out-of-pocket for prescription drugs covered by your plan in 2027.[2]

Important: This payment option does not lower the total cost of your prescriptions, but it might help you manage your monthly expenses. Enrolling earlier in the year gives you more months to spread out your drug costs.[2]

ENROLLING IS EASY

If you have Medicare Part D prescription drug coverage, you can sign up for M3P during Open Enrollment or at any time during the year.

There is no cost to join M3P.[1]

BY PHONE
Call your Medicare health or drug plan to request to participate.

ONLINE
Visit your Medicare health or drug plan website to sign up.

FREQUENTLY ASKED QUESTIONS
Please scan here, or visit:
www.medicare.gov/prescription-payment-plan/before-payment-option

1. https://www.medicare.gov/prescription-payment-plan
2. https://www.medicare.gov/prescription-payment-plan/before-payment-option

Rayaldee®
calcifediol ER capsules
30 mcg

How do you take Rayaldee® (calcifediol) ER capsules?



Take your prescribed dose of Rayaldee **30mcg** by mouth once daily **at bedtime***.

1. Take Rayaldee capsule **whole**. Do not chew, break, or open.
2. If you miss a dose, do not take an extra dose. Simply take Rayaldee® at the next scheduled time.

*Your doctor may increase the dose based on your body's response.

Rayaldee works!

Rayaldee is 25(OH)D in an extended release formulation. It is a treatment for secondary hyperparathyroidism (SHPT) in adults with stage 3 or 4 chronic kidney disease (CKD) and low vitamin D levels (25-hydroxyvitamin D less than 30 ng/mL).

It is not for patients with stage 5 CKD or end-stage kidney disease on dialysis.

Rayaldee helps your body **increase your vitamin D levels** and your parathyroid glands make **less PTH**.

IMPORTANT SAFETY INFORMATION

- Before starting Rayaldee, the level of calcium in your blood should be in the normal range (below 9.8 mg/dL). Your doctor can tell you if your calcium level is right for starting Rayaldee.
- Rayaldee can cause hypercalcemia and hypercalciuria (high levels of calcium in your blood and urine, respectively). You should tell your doctor if you are taking any other medications, including other vitamin D medications (prescription or over-the-counter vitamin D supplements). Your doctor may tell you to stop taking them while you are taking Rayaldee, as very high calcium levels due to high vitamin D levels could cause severe changes to your heart rhythm or seizures and require emergency treatment.
- Tell your doctor about any and all medication you may be taking. While you are taking Rayaldee, tell your doctor if you receive any new prescriptions or start any new over-the-counter medications.
- Tell your doctor immediately if you experience any of the following symptoms of very high calcium: feeling tired, difficulty thinking clearly, loss of appetite, nausea, vomiting, constipation, increased thirst, increased urination, weight loss, or any other abnormal symptoms or side effects.
- While you are taking Rayaldee, your doctor will order laboratory tests to see how your SHPT is doing. These lab tests will also examine your blood levels of calcium, phosphorus, 25-hydroxyvitamin D, and PTH. Based on your test results, your dose of Rayaldee may be changed or stopped until your lab results improve.
- The most common side effects of Rayaldee reported by patients in clinical trials were anemia (low levels of red blood cells or hemoglobin), infection in the nose and/or throat, high levels of blood creatinine (a lab test for measuring kidney function), shortness of breath, cough, congestive heart failure, and constipation.
- Take your Rayaldee at bedtime and swallow your capsules whole; do not bite or chew them.
- If you miss a dose, take Rayaldee at your next scheduled time. Do not take an extra dose to make up for the dose you missed.
- Be sure to follow your doctor's orders, including getting your laboratory tests done on time.

Please see Important Safety Information (ISI) and Full Prescribing Information available at www.Rayaldee.com.

RAYALDEE® is a registered trademark of Eirgen Pharma Ltd. OPKO Renal is a division of OPKO Health, Inc.
© 2026 OPKO Pharmaceuticals, LLC. All rights reserved. OP-US-0931-092026v.2